

# Algebraic Number Theory

(PARI-GP version 2.11.0)

## Binary Quadratic Forms

create  $ax^2 + bxy + cy^2$  (distance  $d$ )      `Qfb( $a, b, c, \{d\}$ )`  
reduce  $x$  ( $s = \sqrt{D}$ ,  $l = \lfloor s \rfloor$ )      `qfbred( $x, \{flag\}, \{D\}, \{l\}, \{s\}$ )`  
return  $[y, g]$ ,  $g \in \text{SL}_2(\mathbf{Z})$ ,  $y = g \cdot x$  reduced      `qfbreds12( $x$ )`  
composition of forms       $x*y$  or `qfbnucomp( $x, y, l$ )`  
 $n$ -th power of form       $x^n$  or `qfbnupow( $x, n$ )`  
composition without reduction      `qfbcomprow( $x, y$ )`  
 $n$ -th power without reduction      `qfbpowrow( $x, n$ )`  
prime form of disc.  $x$  above prime  $p$       `qfbprimeform( $x, p$ )`  
class number of disc.  $x$       `qfbclassno( $x$ )`  
Hurwitz class number of disc.  $x$       `qfbhclassno( $x$ )`  
solve  $Q(x, y) = p$  in integers,  $p$  prime      `qfbsolve( $Q, p$ )`

## Quadratic Fields

quadratic number  $\omega = \sqrt{x}$  or  $(1 + \sqrt{x})/2$       `quadgen( $x$ )`  
minimal polynomial of  $\omega$       `quadpoly( $x$ )`  
discriminant of  $\mathbf{Q}(\sqrt{x})$       `quaddisc( $x$ )`  
regulator of real quadratic field      `quadregulator( $x$ )`  
fundamental unit in real  $\mathbf{Q}(\sqrt{D})$       `quadunit( $D, \{w\}$ )`  
class group of  $\mathbf{Q}(\sqrt{D})$       `quadclassunit( $D, \{flag\}, \{t\}$ )`  
Hilbert class field of  $\mathbf{Q}(\sqrt{D})$       `quadhilbert( $D, \{flag\}$ )`  
... using specific class invariant ( $D < 0$ )      `polclass( $D, \{inv\}$ )`  
ray class field modulo  $f$  of  $\mathbf{Q}(\sqrt{D})$       `quadray( $D, f, \{flag\}$ )`

## General Number Fields: Initializations

The number field  $K = \mathbf{Q}[X]/(f)$  is given by irreducible  $f \in \mathbf{Q}[X]$ . We denote  $\theta = \bar{X}$  the canonical root of  $f$  in  $K$ . A  $nf$  structure contains a maximal order and allows operations on elements and ideals. A  $bnf$  adds class group and units. A  $bnr$  is attached to ray class groups and class field theory. A  $rnf$  is attached to relative extensions  $L/K$ .

init number field structure  $nf$       `nfinit( $f, \{flag\}$ )`  
known integer basis  $B$       `nfinit( $[f, B]$ )`  
order maximal at  $vp = [p_1, \dots, p_k]$       `nfinit( $[f, vp]$ )`  
order maximal at all  $p \leq P$       `nfinit( $[f, P]$ )`  
certify maximal order      `nfcertify( $nf$ )`

### nf members:

a monic  $F \in \mathbf{Z}[X]$  defining  $K$        $nf.pol$   
number of real/complex places       $nf.r1/r2/sign$   
discriminant of  $nf$        $nf.disc$   
 $T_2$  matrix       $nf.t2$   
complex roots of  $F$        $nf.roots$   
integral basis of  $\mathbf{Z}_K$  as powers of  $\theta$        $nf.zk$   
different/codifferent       $nf.diff, nf.codiff$   
index  $[\mathbf{Z}_K : \mathbf{Z}[X]/(F)]$        $nf.index$   
recompute  $nf$  using current precision      `nfnewprec( $nf$ )`  
init relative  $rnf$   $L = K[Y]/(g)$       `rnfinit( $nf, g$ )`  
init  $bnf$  structure      `bnfinit( $f, \{flag\}$ )`  
**bnf members:** same as  $nf$ , plus  
underlying  $nf$        $bnf.nf$   
classgroup       $bnf.clgp$   
regulator       $bnf.reg$   
fundamental/torsion units       $bnf.fu, bnf.tu$

compress a  $bnf$  for storage      `bnfcompress( $bnf$ )`  
recover a  $bnf$  from compressed  $bnfz$       `bnfinit( $bnfz$ )`  
add  $S$ -class group and units, yield  $bnfS$       `bnfsunit( $bnf, S$ )`  
init class field structure  $bnr$       `bnrinit( $bnf, m, \{flag\}$ )`  
**bnr members:** same as  $bnf$ , plus  
underlying  $bnf$        $bnr.bnf$   
big ideal structure       $bnr.bid$   
modulus       $bnr.mod$   
structure of  $(\mathbf{Z}_K/m)^*$        $bnr.zkst$

## Fields, subfields, embeddings

**Defining polynomials, embeddings**  
smallest poly defining  $f = 0$  (slow)      `polredabs( $f, \{flag\}$ )`  
small poly defining  $f = 0$  (fast)      `polredbest( $f, \{flag\}$ )`  
random Tschirnhausen transform of  $f$       `poltschirnhaus( $f$ )`  
 $\mathbf{Q}[t]/(f) \subset \mathbf{Q}[t]/(g)$  ? Isomorphic?      `nfisincl( $f, g$ ), nfisom`  
reverse polmod  $a = A(t) \bmod T(t)$       `modreverse( $a$ )`  
compositum of  $\mathbf{Q}[t]/(f)$ ,  $\mathbf{Q}[t]/(g)$       `polcompositum( $f, g, \{flag\}$ )`  
compositum of  $K[t]/(f)$ ,  $K[t]/(g)$       `nfcompositum( $nf, f, g, \{flag\}$ )`  
splitting field of  $K$  (degree divides  $d$ )      `nfsplitting( $nf, \{d\}$ )`  
signs of real embeddings of  $x$       `nfeltsign( $nf, x, \{pl\}$ )`  
complex embeddings of  $x$       `nfeltembed( $nf, x, \{pl\}$ )`  
 $T \in K[t]$ , # of real roots of  $\sigma(T) \in R[t]$       `nfpolsturm( $nf, T, \{pl\}$ )`

### Subfields, polynomial factorization

subfields (of degree  $d$ ) of  $nf$       `nfsubfields( $nf, \{d\}$ )`  
 $d$ -th degree subfield of  $\mathbf{Q}(\zeta_n)$       `polsubcyclo( $n, d, \{v\}$ )`  
roots of unity in  $nf$       `nfrootsof1( $nf$ )`  
roots of  $g$  belonging to  $nf$       `nfroots( $nf, g$ )`  
factor  $g$  in  $nf$       `nfactor( $nf, g$ )`  
factor  $g \bmod$  prime  $pr$  in  $nf$       `nfactormod( $nf, g, pr$ )`

### Linear and algebraic relations

poly of degree  $\leq k$  with root  $x \in \mathbf{C}$       `algdep( $x, k$ )`  
alg. dep. with pol. coeffs for series  $s$       `seralgdep( $s, x, y$ )`  
small linear rel. on coords of vector  $x$       `lindep( $x$ )`

## Basic Number Field Arithmetic (nf)

Number field elements are `t_INT`, `t_FRAC`, `t_POL`, `t_POLMOD`, or `t_COL` (on integral basis  $nf.zk$ ).

### Basic operations

$x + y$       `nfeltadd( $nf, x, y$ )`  
 $x \times y$       `nfeltmul( $nf, x, y$ )`  
 $x^n$ ,  $n \in \mathbf{Z}$       `nfeltpow( $nf, x, n$ )`  
 $x/y$       `nfeltdiv( $nf, x, y$ )`  
 $q = x \backslash y := \text{round}(x/y)$       `nfeltdiveuc( $nf, x, y$ )`  
 $r = x \% y := x - (x \backslash y)y$       `nfeltmod( $nf, x, y$ )`  
...  $[q, r]$  as above      `nfeltdivrem( $nf, x, y$ )`  
reduce  $x$  modulo ideal  $A$       `nfeltreduce( $nf, x, A$ )`  
absolute trace  $\text{Tr}_{K/\mathbf{Q}}(x)$       `nfelttrace( $nf, x$ )`  
absolute norm  $N_{K/\mathbf{Q}}(x)$       `nfeltnorm( $nf, x$ )`

### Multiplicative structure of $K^*$ ; $K^*/(K^*)^n$

valuation  $vp_{\mathfrak{p}}(x)$       `nfeltval( $nf, x, \mathfrak{p}$ )`  
... write  $x = \pi^{vp_{\mathfrak{p}}(x)}y$       `nfeltval( $nf, x, \mathfrak{p}, \&y$ )`  
quadratic Hilbert symbol (at  $\mathfrak{p}$ )      `nfhilbert( $nf, a, b, \{\mathfrak{p}\}$ )`  
 $b$  such that  $xb^n = v$  is small      `idealredmodpower( $nf, x, n$ )`

### Maximal order and discriminant

integral basis of field  $\mathbf{Q}[x]/(f)$       `nfbasis( $f$ )`  
field discriminant of field  $f = 0$       `nfdisc( $f$ )`  
express  $x$  on integer basis      `nfalgtobasis( $nf, x$ )`  
express element  $x$  as a polmod      `nfbasistoalg( $nf, x$ )`

### Dedekind Zeta Function $\zeta_K$ , Hecke $L$ series

$R = [c, w, h]$  in initialization means we restrict  $s \in \mathbf{C}$  to domain  $|\Re(s) - c| < w$ ,  $|\Im(s)| < h$ ;  $R = [w, h]$  encodes  $[1/2, w, h]$  and  $[h]$  encodes  $R = [1/2, 0, h]$  (critical line up to height  $h$ ).  
 $\zeta_K$  as Dirichlet series,  $N(I) < b$       `dirzetak( $nf, b$ )`  
init  $\zeta_K^{(k)}(s)$  for  $k \leq n$       `L = lfunitinit( $bnf, R, \{n = 0\}$ )`  
compute  $\zeta_K(s)$  ( $n$ -th derivative)      `lfun( $L, s, \{n = 0\}$ )`  
compute  $\Lambda_K(s)$  ( $n$ -th derivative)      `lfunlambda( $L, s, \{n = 0\}$ )`

init  $L_K^{(k)}(s, \chi)$  for  $k \leq n$       `L = lfunitinit( $[bnr, chi], R, \{n = 0\}$ )`  
compute  $L_K(s, \chi)$  ( $n$ -th derivative)      `lfun( $L, s, \{n\}$ )`  
Artin root number of  $K$       `bnrrootnumber( $bnr, chi, \{flag\}$ )`  
 $L(1, \chi)$ , for all  $\chi$  trivial on  $H$       `bnrL1( $bnr, \{H\}, \{flag\}$ )`

## Class Groups & Units (bnf, bnr)

Class field theory data  $a_1, \{a_2\}$  is usually  $bnr$  (ray class field),  $bnr, H$  (congruence subgroup) or  $bnr, \chi$  (character on `bnr.clgp`). Any of these define a unique abelian extension of  $K$ .

remove GRH assumption from  $bnf$       `bnfcertify( $bnf$ )`  
expo. of ideal  $x$  on class gp      `bnfisprincipal( $bnf, x, \{flag\}$ )`  
expo. of ideal  $x$  on ray class gp      `bnrisprincipal( $bnr, x, \{flag\}$ )`  
expo. of  $x$  on fund. units      `bnfisunit( $bnf, x$ )`  
as above for  $S$ -units      `bnfissunit( $bnfs, x$ )`  
signs of real embeddings of  $bnf.fu$       `bnfsignunit( $bnf$ )`  
narrow class group      `bnfnarrow( $bnf$ )`

### Class Field Theory

ray class number for modulus  $m$       `bnrclassno( $bnf, m$ )`  
discriminant of class field      `bnrdisc( $a_1, \{a_2\}$ )`  
ray class numbers,  $l$  list of moduli      `bnrclassnolist( $bnf, l$ )`  
discriminants of class fields      `bnrdisclist( $bnf, l, \{arch\}, \{flag\}$ )`  
decode output from `bnrdisclist`      `bnfdecodemodule( $nf, fa$ )`  
is modulus the conductor?      `bnrconductor( $a_1, \{a_2\}$ )`  
is class field  $(bnr, H)$  Galois over  $K^G$       `bnrisgalois( $bnr, G, H$ )`  
action of automorphism on  $bnr.gen$       `bnrgaloismatrix( $bnr, aut$ )`  
apply `bnrgaloismatrix`  $M$  to  $H$       `bnrgaloisapply( $bnr, M, H$ )`  
characters on `bnr.clgp` s.t.  $\chi(g_i) = e(v_i)$       `bnrchar( $bnr, g, \{v\}$ )`  
conductor of character  $\chi$       `bnrconductor( $bnr, chi$ )`  
conductor of extension      `bnrconductor( $a_1, \{a_2\}, \{flag\}$ )`  
conductor of extension  $K[Y]/(g)$       `rnfconductor( $bnf, g$ )`  
Artin group of extension  $K[Y]/(g)$       `rnfnormgroup( $bnr, g$ )`  
subgroups of  $bnr$ , index  $\leq b$       `subgrouplist( $bnr, b, \{flag\}$ )`  
rel. eq. for class field def'd by  $sub$       `rnfkummer( $bnr, sub, \{d\}$ )`  
same, using Stark units (real field)      `bnrstark( $bnr, sub, \{flag\}$ )`  
is  $a$  an  $n$ -th power in  $K_v$  ?      `nfislocalpower( $nf, v, a, n$ )`  
cyclic  $L/K$  satisf. local conditions      `nfgrunwaldwang( $nf, P, D, pl$ )`

### Logarithmic class group

logarithmic  $\ell$ -class group      `bnflog( $bnf, \ell$ )`  
 $[\bar{e}(F_v/Q_p), \bar{f}(F_v/Q_p)]$       `bnflogef( $bnf, pr$ )`  
 $\exp \deg_F(A)$       `bnflogdegree( $bnf, A, \ell$ )`  
is  $\ell$ -extension  $L/K$  locally cyclotomic      `rnfislocalcyclo( $rnf$ )`

**Ideals:** elements, primes, or matrix of generators in HNF

is  $id$  an ideal in  $nf$  ? nfsideal( $nf, id$ )  
is  $x$  principal in  $bnf$  ? bnfisprincipal( $bnf, x$ )  
give  $[a, b]$ , s.t.  $a\mathbf{Z}_K + b\mathbf{Z}_K = x$  idealtwoelt( $nf, x, \{a\}$ )  
put ideal  $a$  ( $a\mathbf{Z}_K + b\mathbf{Z}_K$ ) in HNF form idealhnf( $nf, a, \{b\}$ )  
norm of ideal  $x$  idealnrm( $nf, x$ )  
minimum of ideal  $x$  (direction  $v$ ) idealmin( $nf, x, v$ )  
LLL-reduce the ideal  $x$  (direction  $v$ ) idealred( $nf, x, \{v\}$ )

**Ideal Operations**

add ideals  $x$  and  $y$  idealadd( $nf, x, y$ )  
multiply ideals  $x$  and  $y$  idealmul( $nf, x, y, \{flag\}$ )  
intersection of ideals  $x$  and  $y$  idealintersect( $nf, x, y, \{flag\}$ )  
 $n$ -th power of ideal  $x$  idealpow( $nf, x, n, \{flag\}$ )  
inverse of ideal  $x$  idealinv( $nf, x$ )  
divide ideal  $x$  by  $y$  idealdiv( $nf, x, y, \{flag\}$ )  
Find  $(a, b) \in x \times y, a + b = 1$  idealaddtoone( $nf, x, \{y\}$ )  
coprime integral  $A, B$  such that  $x = A/B$  idealnumden( $nf, x$ )

**Primes and Multiplicative Structure**

factor ideal  $x$  in  $\mathbf{Z}_K$  idealfactor( $nf, x$ )  
expand ideal factorization in  $K$  idealfactorback( $nf, f, \{e\}$ )  
is ideal  $A$  an  $n$ -th power ? idealispower( $nf, A, n$ )  
expand elt factorization in  $K$  nffactorback( $nf, f, \{e\}$ )  
decomposition of prime  $p$  in  $\mathbf{Z}_K$  idealprimedec( $nf, p$ )  
valuation of  $x$  at prime ideal  $pr$  idealval( $nf, x, pr$ )  
weak approximation theorem in  $nf$  idealchinese( $nf, x, y$ )  
 $a \in K$ , s.t.  $v_p(a) = v_p(x)$  if  $v_p(x) \neq 0$  idealappr( $nf, x$ )  
 $a \in K$  such that  $(a \cdot x, y) = 1$  idealcoprime( $nf, x, y$ )  
give  $bid$  =structure of  $(\mathbf{Z}_K/id)^*$  idealstar( $nf, id, \{flag\}$ )  
structure of  $(1 + \mathfrak{p})/(1 + \mathfrak{p}^k)$  idealprincipalunits( $nf, pr, k$ )  
discrete log of  $x$  in  $(\mathbf{Z}_K/bid)^*$  ideallog( $nf, x, bid$ )  
**idealstar** of all ideals of norm  $\leq b$  ideallist( $nf, b, \{flag\}$ )  
add Archimedean places ideallistarch( $nf, b, \{ar\}, \{flag\}$ )  
init **modpr** structure nfmodprinit( $nf, pr$ )  
project  $t$  to  $\mathbf{Z}_K/pr$  nfmodpr( $nf, t, modpr$ )  
lift from  $\mathbf{Z}_K/pr$  nfmodprlift( $nf, t, modpr$ )

**Galois theory over  $\mathbf{Q}$**

conjugates of a root  $\theta$  of  $nf$  nfgaloisconj( $nf, \{flag\}$ )  
apply Galois automorphism  $s$  to  $x$  nfgaloisapply( $nf, s, x$ )  
Galois group of field  $\mathbf{Q}[x]/(f)$  polgalois( $f$ )  
initializes a Galois group structure  $G$  galoisinit( $pol, \{den\}$ )  
character table of  $G$  galoischartable( $G$ )  
conjugacy classes of  $G$  galoisconjclasses( $G$ )  
 $\det(1 - \rho(g)T)$ ,  $\chi$  character of  $\rho$  galoischarpoly( $G, \chi, \{o\}$ )  
 $\det(\rho(g))$ ,  $\chi$  character of  $\rho$  galoischarDET( $G, \chi, \{o\}$ )  
action of  $p$  in nfgaloisconj form galoispermtopol( $G, \{p\}$ )  
identify as abstract group galoisidentify( $G$ )  
export a group for GAP/MAGMA galoisexport( $G, \{flag\}$ )  
subgroups of the Galois group  $G$  galoissubgroups( $G$ )  
is subgroup  $H$  normal? galoisisnormal( $G, H$ )  
subfields from subgroups galoissubfields( $G, \{flag\}, \{v\}$ )  
fixed field galoisfixedfield( $G, perm, \{flag\}, \{v\}$ )  
Frobenius at maximal ideal  $P$  idealfrobenius( $nf, G, P$ )  
ramification groups at  $P$  idealramgroups( $nf, G, P$ )  
is  $G$  abelian? galoisisabelian( $G, \{flag\}$ )  
abelian number fields/ $\mathbf{Q}$  galoissubcyclo( $N, H, \{flag\}, \{v\}$ )

**Algebraic Number Theory**

(PARI-GP version 2.11.0)

**The galpol package**

query the package: polynomial galoisgetpol(a,b,{s})  
... : permutation group galoisgetgroup(a,b)  
... : group description galoisgetname(a,b)

**Relative Number Fields (rnf)**

Extension  $L/K$  is defined by  $T \in K[x]$ . rnfequation( $nf, T, \{flag\}$ )  
absolute equation of  $L$  rnfisabelian( $nf, T$ )  
is  $L/K$  abelian? rnfalgtobasis( $rnf, x$ )  
relative nfalgtobasis rnfbasistoalg( $rnf, x$ )  
relative nfbasistoalg rnfidealhnf( $rnf, x$ )  
relative idealhnf rnfidealmul( $rnf, x, y$ )  
relative idealmul rnfidealtwoelt( $rnf, x$ )  
relative idealtwoelt

**Lifts and Push-downs**

absolute  $\rightarrow$  relative representation for  $x$  rnfeltabstorel( $rnf, x$ )  
relative  $\rightarrow$  absolute representation for  $x$  rnfeltreltoabs( $rnf, x$ )  
lift  $x$  to the relative field rnfeltup( $rnf, x$ )  
push  $x$  down to the base field rnfeltdown( $rnf, x$ )  
idem for  $x$  ideal: (rnfideal)reltoabs, abstorel, up, down

**Norms and Trace**

relative norm of element  $x \in L$  rnfeltnrm( $rnf, x$ )  
relative trace of element  $x \in L$  rnfelttrace( $rnf, x$ )  
absolute norm of ideal  $x$  rnfidealnrmabs( $rnf, x$ )  
relative norm of ideal  $x$  rnfidealnrmrel( $rnf, x$ )  
solutions of  $N_{K/\mathbf{Q}}(y) = x \in \mathbf{Z}$  bnfisintnorm( $bnf, x$ )  
is  $x \in \mathbf{Q}$  a norm from  $K$ ? bnfisnorm( $bnf, x, \{flag\}$ )  
initialize  $T$  for norm eq. solver rnfisnorminit( $K, pol, \{flag\}$ )  
is  $a \in K$  a norm from  $L$ ? rnfisnorm( $T, a, \{flag\}$ )  
initialize  $t$  for Thue equation solver thueinit( $f$ )  
solve Thue equation  $f(x, y) = a$  thue( $t, a, \{sol\}$ )  
characteristic poly. of  $a$  mod  $T$  rnfcharpoly( $nf, T, a, \{v\}$ )

**Factorization**

factor ideal  $x$  in  $L$  rnfidealfactor( $rnf, x$ )  
 $[S, T]: T_{i,j} \mid S_i; S$  primes of  $K$  above  $p$  rnfidealprimedec( $rnf, p$ )

**Maximal order  $\mathbf{Z}_L$  as a  $\mathbf{Z}_K$ -module**

relative polredbest rnfpolredbest( $nf, T$ )  
relative polredabs rnfpolredabs( $nf, T$ )  
relative Dedekind criterion, prime  $pr$  rnfdedekind( $nf, T, pr$ )  
discriminant of relative extension rnfdisc( $nf, T$ )  
pseudo-basis of  $\mathbf{Z}_L$  rnfpsedobasis( $nf, T$ )

**General  $\mathbf{Z}_K$ -modules:**  $M = [\text{matrix, vec. of ideals}] \subset L$   
relative HNF / SNF nfhnf( $nf, M$ ), nfsnf  
multiple of det  $M$  nfdetint( $nf, M$ )  
HNF of  $M$  where  $d = nfdetint(M)$  nfhnfmod( $x, d$ )  
reduced basis for  $M$  rnflllgram( $nf, T, M$ )  
determinant of pseudo-matrix  $M$  rnfDET( $nf, M$ )  
Steinitz class of  $M$  rnfsteinitz( $nf, M$ )  
 $\mathbf{Z}_K$ -basis of  $M$  if  $\mathbf{Z}_K$ -free, or 0 rnfhnfbasis( $bnf, M$ )  
 $n$ -basis of  $M$ , or  $(n + 1)$ -generating set rnfBasis( $bnf, M$ )  
is  $M$  a free  $\mathbf{Z}_K$ -module? rnfisfree( $bnf, M$ )

**Associative Algebras**

$A$  is a general associative algebra given by a multiplication table  $mt$  (over  $\mathbf{Q}$  or  $\mathbf{F}_p$ ); represented by  $al$  from **algtabinit**.  
create  $al$  from  $mt$  (over  $\mathbf{F}_p$ ) algtabinit( $mt, \{p = 0\}$ )  
group algebra  $\mathbf{Q}[G]$  (or  $\mathbf{F}_p[G]$ ) alggroup( $G, \{p = 0\}$ )  
center of group algebra alggroupcenter( $G, \{p = 0\}$ )

**Properties**

is  $(mt, p)$  OK for **algtabinit**? algisassociative( $mt, \{p = 0\}$ )  
multiplication table  $mt$  algmtable( $al$ )  
dimension of  $A$  over prime subfield algdim( $al$ )  
characteristic of  $A$  algchar( $al$ )  
is  $A$  commutative? algiscommutative( $al$ )  
is  $A$  simple? algissimple( $al$ )  
is  $A$  semi-simple? algissemisimple( $al$ )  
center of  $A$  algcenter( $al$ )  
Jacobson radical of  $A$  algradical( $al$ )  
radical  $J$  and simple factors of  $A/J$  algsimpledec( $al$ )

**Operations on algebras**

create  $A/I, I$  two-sided ideal algquotient( $al, I$ )  
create  $A_1 \otimes A_2$  algtensor( $al1, al2$ )  
create subalgebra from basis  $B$  algsubalg( $al, B$ )  
quotients by ortho. central idempotents  $e$  algcentralproj( $al, e$ )  
isomorphic alg. with integral mult. table algmakeintegral( $mt$ )  
prime subalgebra of semi-simple  $A$  over  $\mathbf{F}_p$  algprimesubalg( $al$ )  
find isomorphism  $A \cong M_d(\mathbf{F}_q)$  algsplit( $al$ )

**Operations on lattices in algebras**

lattice generated by cols. of  $M$  alglathnf( $al, M$ )  
... by the products  $xy, x \in lat1, y \in lat2$  alglatmul( $al, lat1, lat2$ )  
sum  $lat1 + lat2$  of the lattices alglatadd( $al, lat1, lat2$ )  
intersection  $lat1 \cap lat2$  alglatinter( $al, lat1, lat2$ )  
test  $lat1 \subset lat2$  alglatsubset( $al, lat1, lat2$ )  
generalized index  $(lat2 : lat1)$  alglatindex( $al, lat1, lat2$ )  
 $\{x \in al \mid x \cdot lat1 \subset lat2\}$  alglatlefttransporter( $al, lat1, lat2$ )  
 $\{x \in al \mid lat1 \cdot x \subset lat2\}$  alglatrightrighttransporter( $al, lat1, lat2$ )  
test  $x \in lat$  (set  $c = \text{coord. of } x$ ) alglatcontains( $al, lat, x, \{&c\}$ )  
element of  $lat$  with coordinates  $c$  alglatelement( $al, lat, c$ )

**Operations on elements**

$a + b, a - b, -a$  algadd( $al, a, b$ ), algsub, algneg  
 $a \times b, a^2$  algmul( $al, a, b$ ), algsqr  
 $a^n, a^{-1}$  algpow( $al, a, n$ ), alginv  
is  $x$  invertible ? (then set  $z = x^{-1}$ ) alginv( $al, x, \{&z\}$ )  
find  $z$  such that  $x \times z = y$  algdivl( $al, x, y$ )  
find  $z$  such that  $z \times x = y$  algdivr( $al, x, y$ )  
does  $z$  s.t.  $x \times z = y$  exist? (set it) algisdivl( $al, x, y, \{&z\}$ )  
matrix of  $v \mapsto x \cdot v$  algtomatrix( $al, x$ )  
absolute norm algnorm( $al, x$ )  
absolute trace algtrace( $al, x$ )  
absolute char. polynomial algcharpoly( $al, x$ )  
given  $a \in A$  and polynomial  $T$ , return  $T(a)$  algpoleval( $al, T, a$ )  
random element in a box algrandom( $al, b$ )

Based on an earlier version by Joseph H. Silverman

July 2018 v2.35. Copyright © 2018 K. Belabas

Permission is granted to make and distribute copies of this card provided the copyright and this permission notice are preserved on all copies.

Send comments and corrections to (Karim.Belabas@math.u-bordeaux.fr)

Central Simple Algebras

$A$  is a central simple algebra over a number field  $K$ ; represented by  $al$  from **alginit**;  $K$  is given by a  $nf$  structure.

create CSA from data	<b>alginit</b> ( $B, C, \{v\}, \{maxord = 1\}$ )
multiplication table over $K$	$B = K, C = mt$
cyclic algebra ( $L/K, \sigma, b$ )	$B = rnf, C = [sigma, b]$
quaternion algebra $(a, b)_K$	$B = K, C = [a, b]$
matrix algebra $M_d(K)$	$B = K, C = d$
local Hasse invariants over $K$	$B = K, C = [d, [PR, HF], HI]$

Properties

type of $al$ ( $mt, CSA$ )	<b>algtype</b> ( $al$ )
dimension of $A$ over $\mathbf{Q}$	<b>algdim</b> ( $al, 1$ )
dimension of $al$ over its center $K$	<b>algdim</b> ( $al$ )
degree of $A$ ( $= \sqrt{\dim_K A}$ )	<b>algdegree</b> ( $al$ )
$al$ a cyclic algebra ( $L/K, \sigma, b$ ); return $\sigma$	<b>algaut</b> ( $al$ )
...return $b$	<b>algb</b> ( $al$ )
...return $L/K$ , as an $rnf$	<b>algsplittingfield</b> ( $al$ )
split $A$ over an extension of $K$	<b>algsplittingdata</b> ( $al$ )
splitting field of $A$ as an $rnf$ over center	<b>algsplittingfield</b> ( $al$ )
multiplication table over center	<b>algrelmultable</b> ( $al$ )
places of $K$ at which $A$ ramifies	<b>algramifiedplaces</b> ( $al$ )
Hasse invariants at finite places of $K$	<b>alghassef</b> ( $al$ )
Hasse invariants at infinite places of $K$	<b>alghassei</b> ( $al$ )
Hasse invariant at place $v$	<b>alghasse</b> ( $al, v$ )
index of $A$ over $K$ (at place $v$ )	<b>algindex</b> ( $al, \{v\}$ )
is $al$ a division algebra? (at place $v$ )	<b>algisdivision</b> ( $al, \{v\}$ )
is $A$ ramified? (at place $v$ )	<b>algisramified</b> ( $al, \{v\}$ )
is $A$ split? (at place $v$ )	<b>algissplit</b> ( $al, \{v\}$ )

Operations on elements

reduced norm	<b>algnorm</b> ( $al, x$ )
reduced trace	<b>algtrace</b> ( $al, x$ )
reduced char. polynomial	<b>algcharpoly</b> ( $al, x$ )
express $x$ on integral basis	<b>algalgtobasis</b> ( $al, x$ )
convert $x$ to algebraic form	<b>algbasistoalg</b> ( $al, x$ )
map $x \in A$ to $M_d(L)$ , $L$ split. field	<b>algtomatrix</b> ( $al, x$ )

Orders

<b>Z</b> -basis of order $\mathcal{O}_0$	<b>algbasis</b> ( $al$ )
discriminant of order $\mathcal{O}_0$	<b>algdisc</b> ( $al$ )
<b>Z</b> -basis of natural order in terms $\mathcal{O}_0$ 's basis	<b>alginvbasis</b> ( $al$ )